arXiv:1411.7317v2 [math-ph] 27 Nov 2014

Theory of Classical Higgs Fields. II. Lagrangians

G. SARDANASHVILY, A. KUROV

Department of Theoretical Physics, Moscow State University, Russia

Abstract

We consider classical gauge theory with spontaneous symmetry breaking on a principal bundle $P \to X$ whose structure group G is reducible to a closed subgroup H, and sections of the quotient bundle $P/H \to X$ are treated as classical Higgs fields. In this theory, matter fields with an exact symmetry group H are described by sections of a composite bundle $Y \to P/H \to X$. We show that their gauge G-invariant Lagrangian necessarily factorizes through a vertical covariant differential on Y defined by a principal connection on an H-principal bundle $P \to P/H$ (Theorems 5 and 6).

Following our previous work [10], we consider classical gauge theory on a principal bundle $P \to X$ with a structure Lie group G which is reducible to its closed subgroup H, i.e., P admits reduced principal subbundles possessing a structure group H.

Given a closed (and, consequently, Lie) subgroup $H \subset G$, we have a composite bundle

$$P \to P/H \to X, \tag{1}$$

where

$$P_\Sigma = P \longrightarrow P/H \tag{2}$$

is a principal bundle with a structure group H and

$$\Sigma = P/H \longrightarrow X \tag{3}$$

is a P-associated bundle with a typical fibre G/H which a structure group G acts on by left multiplications. In accordance with the well-known theorem [1, 12], there is one-to-one correspondence between the global sections h of the quotient bundle (3) and the reduced H-principal subbundles P^h of P which are the restriction

$$P^h = h^* P_\Sigma \tag{4}$$

of the H-principal bundle P_Σ (2) to $h(X) \subset \Sigma$. In classical gauge theory, global sections of the quotient bundle (3) are treated as classical Higgs fields [1, 8, 11].

A question is how to describe matter fields in gauge theory with a structure group G if they admit only an exact symmetry subgroup H. In particular, this is the case of spinor fields in gravitation theory [2, 9].

We have shown that such matter fields are represented by sections of a composite bundle

$$\pi_{YX} : Y \longrightarrow \Sigma \longrightarrow X \tag{5}$$

where $Y \to \Sigma$ is a P_Σ-associated bundle

$$Y = (P \times V)/H \tag{6}$$

with a structure group H acting on its typical fibre V on the left [1, 8, 11]. Given a global section h of the fibre bundle $\Sigma \to X$ (3), the restriction

$$Y^h = h^*Y = (h^*P \times V)/H = (P^h \times V)/H \tag{7}$$

of a fibre bundle $Y \to \Sigma$ to $h(X) \subset \Sigma$ is a fibre bundle associated with the reduced H-principal subbundle P^h (4) of a G-principal bundle P. Sections of the fibre bundle $Y^h \to X$ (7) describe matter fields in the presence of a Higgs field h. A key point is that the composite bundle $\pi_{YX} : Y \to X$ (5) is proved to be a P-associated bundle

$$Y = (P \times (G \times V)/H)/G$$

with a structure group G [1, 10, 11]. Its typical fibre is a fibre bundle

$$\pi_{WH} : W = (G \times V)/H \to G/H \tag{8}$$

associated with an H-principal bundle $G \to G/H$. A structure group G acts on the W (8) by the induced representation [5]:

$$g : (G \times V)/H \to (gG \times V)/H. \tag{9}$$

This fact enables one to describe matter fields with an exact symmetry group $H \subset G$ in the framework of gauge theory on a G-principal bundle $P \to X$ if its structure group G is reducible to H. Here, we aim to show that their gauge G-invariant Lagrangian necessarily factorizes through a vertical covariant differential on Y defined by an H-principal connection on $P \to P/H$ (Theorems 5 and 6).

A problem is that, though the P-associated composite bundle $Y \to X$ (5 can be endowed with a principal connection on a G-principal bundle $P \to X$, such a connection need not be reducible to principal connections on reduced H-principal subbundles P^h, unless the following condition holds [1, 3].

Theorem 1. *Let a Lie algebra $\mathfrak{g}$ of G be a direct sum*

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{f} \tag{10}$$

of a Lie algebra $\mathfrak{h}$ of H and its supplement $\mathfrak{f}$ obeying the commutation relations

$$[\mathfrak{f}, \mathfrak{f}] \subset \mathfrak{h}, \qquad [\mathfrak{f}, \mathfrak{h}] \subset \mathfrak{f}.$$

(e.g., H is a Cartan subgroup of G). Let A be a principal connection on P. The $\mathfrak{h}$-valued component $\overline{A}^h$ of its pull-back onto a reduced H-principal subbundle P^h is a principal connection on P^h.

At the same time, connections on reduced H-principal subbundles P^h can be generated in a different way. Let $\Pi \to Z$ be a principal bundle with a structure group K. Given a manifold map $\phi : Z' \to Z$, a pull-back bundle $\phi^*\Pi \to Z'$ also is a principal bundle with a structure group K. Let A be a principal connection on a principal bundle $\Pi \to Z$. Then the pull-back connection ϕ^*A is a principal connection on $\phi^*\Pi \to Z'$ [3]. The following is a corollary of this fact.

Theorem 2. *Given the composite bundle (1), let A_Σ be a principal connection on the H-principal bundle $P \to \Sigma$ (2). Then, for any reduced H principal subbundle P^h (4) the pull-back connection h^*A_Σ is a principal connection on P^h.*

Turn now to the composite bundle Y (5). Given an atlas Ψ_P of P, the associated quotient bundle $\Sigma \to X$ (3) is provided with bundle coordinates (x^λ, σ^m). With this atlas and an atlas $\Psi_{Y\Sigma}$ of $Y \to \Sigma$, the composite bundle $Y \to X$ (5) is endowed with adapted bundle coordinates $(x^\lambda, \sigma^m, y^i)$ where (y^i) are fibre coordinates on $Y \to \Sigma$. Then the following holds [1, 8, 11].

Theorem 3. *Let*

$$A_\Sigma = dx^\lambda \otimes (\partial_\lambda + A^a_\lambda e_a) + d\sigma^m \otimes (\partial_m + A^a_m e_a) \tag{11}$$

be a principal connection on an H-principal bundle $P \to \Sigma$ where $\{e_a\}$ is a basis for a Lie algebra $\mathfrak{h}$ of H. Let

$$A_{Y\Sigma} = dx^\lambda \otimes (\partial_\lambda + A^a_\lambda(x^\mu, \sigma^k) I^i_a \partial_i) + d\sigma^m \otimes (\partial_m + A^a_m(x^\mu, \sigma^k) I^i_a \partial_i) \tag{12}$$

be an associated principal connection on $Y \to \Sigma$ where $\{I_a\}$ is a representation of a Lie algebra $\mathfrak{h}$ in V. Then, for any subbundle $Y^h \to X$ (7) of a composite bundle $Y \to X$, the pull-back connection

$$A_h = h^* A_{Y\Sigma} = dx^\lambda \otimes [\partial_\lambda + (A^a_m(x^\mu, h^k)\partial_\lambda h^m + A^a_\lambda(x^\mu, h^k)) I^i_a \partial_i], \tag{13}$$

is a connection on Y^h associated with the pull-back principal connection $h^* A_\Sigma$ on the reduced H-principal subbundle P^h in Theorem 2.

Any connection A_Σ (11) on a fibre bundle $Y \to \Sigma$ yields a first order differential operator, called the vertical covariant differential,

$$\widetilde{D} : J^1Y \to T^*X \underset{Y}{\otimes} V_\Sigma Y, \qquad \widetilde{D} = dx^\lambda \otimes (y^i_\lambda - A^i_\lambda - A^i_m \sigma^m_\lambda)\partial_i, \tag{14}$$

on a composite bundle $Y \to X$ where $V_\Sigma Y$ is the vertical tangent bundle of $Y \to \Sigma$. It possesses the following important property [1, 8, 11].

Theorem 4. *For any section h of a fibre bundle $\Sigma \to X$, the restriction of the vertical differential $\widetilde{D}$ (14) onto the fibre bundle Y^h (7) coincides with a covariant differential D^{A_h} defined by the connection A_h (13) on Y^h.*

In view of Theorems 3 and 4, one can assume that a Lagrangian of matter fields represented by sections of the composite bundle (5) factorizes through the vertical covariant differential $\widetilde{D}$ (14) of some connection $A_{Y\Sigma}$ on a fibre bundle $Y \to \Sigma$. Forthcoming Theorem 5 shows that this factorization is necessary in order that a matter field Lagrangian to be gauge invariant.

Theorem 5. *In gauge theory on a principal bundle P whose structure group G is reducible to a closed subgroup H, a matter field Lagrangian is gauge invariant only if it factorizes through a vertical covariant differential of some principal connection on the H-principal bundle $P \to P/H$ (2).*

Proof. Let $P \to X$ be a principal bundle whose structure group G is reducible to a closed subgroup H. Let Y be the P_Σ-associated bundle (6). A total configuration space of gauge theory of principal connections on P in the presence of matter and Higgs fields is

$$J^1C \underset{X}{\times} J^1Y \tag{15}$$

where $C = J^1P/G$ is the bundle of principal connections on P and J^1Y is the first order jet manifold of $Y \to X$. A total Lagrangian on the configuration space (15) is a sum

$$L_{\rm tot} = L_{\rm A} + L_{\rm m} + L_\sigma \tag{16}$$

of a gauge field Lagrangian $L_{\rm A}$, a matter field Lagrangian $L_{\rm m}$ and a Higgs field Lagrangian L_σ. The total Lagrangian $L_{\rm tot}$ (16) is required to be invariant with respect to vertical principal automorphisms of a G-principal bundle $P \to X$. Any vertical principal automorphism of a G-principal bundle $P \to X$, being G-equivariant, also is H-equivariant and, thus, it is a principal automorphism of an H-principal bundle $P \to \Sigma$. Consequently, it yields an automorphism of the P_Σ-associated bundle Y (5). Accordingly, every G-principal vector field ξ on $P \to X$ (an infinitesimal generator of a local one-parameter group of vertical principal automorphisms of P) also is an H-principal vector field on $P \to \Sigma$. It yields an infinitesimal gauge transformation υ_ξ of a composite bundle Y seen as a P- and P_Σ-associated bundle. This reads

$$\upsilon_\xi = \xi^p(x^\mu)J^m_p\partial_m + \vartheta^a_\xi(x^\mu,\sigma^k)I^A_a\partial_A, \tag{17}$$

where $\{J_p\}$ is a representation of a Lie algebra $\mathfrak{g}$ of G in G/H and $\{I_a\}$ is a representation of a Lie algebra $\mathfrak{h}$ of H in V. Since gauge and Higgs field Lagrangians in the absence of matter fields are assumed to be gauge invariant, a matter field Lagrangian $L_{\rm m}$ also is separately gauge invariant. This means that its Lie derivative along the jet prolongation $J^1\upsilon_\xi$ of the vector field υ_ξ (17) vanishes, that is,

$$\mathbf{L}_{J^1\upsilon_\xi}L_{\rm m} = 0. \tag{18}$$

In order to satisfy the conditions (18), let us consider some principal connection A_Σ (11) on an H-principal bundle $P \to \Sigma$ and the associated connection $A_{Y\Sigma}$ (12) on $Y \to \Sigma$. Let a matter field Lagrangian $L_{\rm m}$ factorize as

$$L_{\rm m} : J^1Y \stackrel{\widetilde{D}}{\to} T^*X \underset{Y}{\otimes} V_\Sigma Y \to \stackrel{n}{\wedge} T^*X$$

through the vertical covariant differential $\widetilde{D}$ (14). In this case, $L_{\rm m}$ can be regarded as a function $L_{\rm m}(y^i, k^i_\lambda)$ of formal variables y^i and $k^i_\lambda = \widetilde{D}^i_\lambda$. The corresponding infinitesimal gauge transformation of variables (y^i, k^i_λ) reads

$$\upsilon = \upsilon^i\partial_i + \partial_j\upsilon^i k^j_\lambda\frac{\partial}{\partial k^i_\lambda}.$$

It is independent of derivatives of gauge parameters ξ. Therefore, the gauge invariance condition is trivially satisfied. □

However, a problem is that the principal connection A_Σ (11) on an H-principal bundle $P \to P/H$ fails to be a dynamic variable in gauge theory. Therefore, let us assume that a Lie algebra of a structure group G satisfies the decomposition (10). In this case, any G-principal connection A on a principal bundle $P \to X$ yields H-principal connections on reduced H-principal subbundles P^h in accordance with Theorem 1. Then one can state the following [11].

Theorem 6. *There exists a connection $A_{Y\Sigma}$ (12) on a fibre bundle $Y \to P/H$ whose restriction $A_h = h^*A_\Sigma$ onto a P^h-associated bundle Y^h coincides with a principal connection $\overline{A}_h$ on P^h generated by a principal connection A on a principal bundle $P \to X$.*

Proof. Let $P^h \subset P$ be a reduced principal subbundle and $\overline{A}_h$ an H-principal connection on P^h in Theorem 1 which is generated by a G-principal connection A on a principal bundle $P \to X$. It is extended to a G-principal connection on P so that h is an integral section of the associated connection

$$\overline{A}_h = dx^\lambda \otimes (\partial_\lambda + A^p_\lambda J^m_p \partial_m)$$

on a P-associated bundle $\Sigma \to X$. Let $\Psi_{Y\Sigma}$ be an atlas of a P_Σ-associated bundle $Y \to \Sigma$ which is defined by a family $\{z_\iota\}$ of local sections of $P \to \Sigma$. Given a section h of $\Sigma \to X$, we have a family of sections $\{z_\iota \circ h\}$ which yields an atlas Ψ^h of a principal bundle $P \to X$ with H-valued transition functions. With respect to this atlas, a section h takes its values in the center of a quotient space G/H and the connection $\overline{A}_h$ reads

$$\overline{A}_h = dx^\lambda \otimes (\partial_\lambda + A^a_\lambda e_a). \tag{19}$$

We have

$$A = \overline{A}_h + \Theta = dx^\lambda \otimes (\partial_\lambda + A^a_\lambda e_a) + \Theta^b_\lambda dx^\lambda \otimes e_b, \tag{20}$$

where $\{e_a\}$ is a basis for the Lie algebra $\mathfrak{h}$ and $\{e_b\}$ is that for $\mathfrak{m}$. Written with respect to an arbitrary atlas of P, the decomposition (20) reads

$$A = \overline{A}_h + \Theta, \qquad \Theta = \Theta^p_\lambda dx^\lambda \otimes e_p,$$

and obeys the relation

$$\Theta^p_\lambda J^m_p = \nabla^A_\lambda h^m,$$

where D_λ are covariant derivatives relative to the associated principal connection A on $\Sigma \to X$. Based on this fact, let consider the covariant differential

$$D = D^m_\lambda dx^\lambda \otimes \partial_m = (\sigma^m_\lambda - A^p_\lambda J^m_p) dx^\lambda \otimes \partial_m$$

relative to the associated principal connection A on $\Sigma \to X$. It can be regarded as a $V\Sigma$-valued one-form on the jet manifold $J^1\Sigma$ of $\Sigma \to X$. Since the decomposition (20) holds for any section h of $\Sigma \to X$, there exists a (VP/P)-valued (where VP is the vertical tangent bundle of $P \to X$) one-form

$$\Theta = \Theta^p_\lambda dx^\lambda \otimes e_p$$

on $J^1\Sigma$ which obeys the equation

$$\Theta^p_\lambda J^m_p = D^m_\lambda. \tag{21}$$

Then we obtain the (VP/G)-valued one-form

$$A_H = dx^\lambda \otimes (\partial_\lambda + (A^p_\lambda - \Theta^p_\lambda) e_p)$$

on $J^1\Sigma$ whose pull-back onto each $J^1h(X) \subset J^1\Sigma$ is the connection $\overline{A}_h$ (19) written with respect to the atlas Ψ^h. The decomposition (20) holds and, consequently, the equation (21) possesses a solution for each principal connection A. Therefore, there exists a $(VP)/G$-valued one-form

$$A_H = dx^\lambda \otimes (\partial_\lambda + (a^p_\lambda - \Theta^p_\lambda) e_p) \tag{22}$$

on the product $J^1\Sigma \underset{X}{\times} J^1C$ such that, for any principal connection A and any Higgs field h, the restriction of A_H (22) to

$$J^1h(X) \times A(X) \subset J^1\Sigma \underset{X}{\times} J^1C$$

is the connection $\overline{A}_h$ (19) written with respect to the atlas Ψ^h. Let us now assume that, whenever A is a principal connection on a G-principal bundle $P \to X$, there exists a principal connection A_Σ (11) on a principal H-bundle $P \to \Sigma$ such that the pull-back connection $A_h = h^*A_{Y\Sigma}$ (13) on Y^h coincides with $\overline{A}_h$ (19) for any $h \in \Sigma(X)$. In this case, there exists $V_\Sigma Y$-valued one-form

$$\widetilde{\mathcal{D}} = dx^\lambda \otimes (y^i_\lambda - (A^a_m\sigma^m_\lambda + A^a_\lambda)I^i_a)\partial_i \tag{23}$$

on the configuration space (15) whose components are defined as follows. Given a point

$$(x^\lambda, a^r_\mu, a^r_{\lambda\mu}, \sigma^m, \sigma^m_\lambda, y^i, y^i_\lambda) \in J^1C \underset{X}{\times} J^1Y, \tag{24}$$

let h be a section of $\Sigma \to X$ whose first jet $j^1_x h$ at $x \in X$ is $(\sigma^m, \sigma^m_\lambda)$, i.e.,

$$h^m(x) = \sigma^m, \qquad \partial_\lambda h^m(x) = \sigma^m_\lambda.$$

Let the bundle of principal connections C and the Lie algebra bundle VP/G be provided with the atlases associated with the above mentioned atlas Ψ^h. Then we write

$$A_h = \overline{A}_h, \qquad A^a_m\sigma^m_\lambda + A^a_\lambda = a^a_\lambda - \Theta^a_\lambda. \tag{25}$$

These equations for functions A^a_m and A^a_λ at the point (24) have a solution because Θ^a_λ are affine functions in the jet coordinates σ^m_λ. □

Given solutions of the equations (25) at all points of the configuration space (15), we require that a matter field Lagrangian factorizes as

$$L_{\rm m} : J^1C \underset{X}{\times} J^1Y \overset{\widetilde{D}}{\to} T^*X \underset{Y}{\otimes} V_\Sigma Y \to \overset{n}{\wedge} T^*X \tag{26}$$

through the form $\widetilde{D}$ (23). As a result, we obtain a gauge theory of gauge potentials of a group G, matter fields with an exact symmetry subgroup $H \subset G$ and classical Higgs fields on the configuration space (24).

As was mentioned above, an example of classical Higgs fields is a metric gravitational field in gauge gravitation theory on natural bundles with spontaneous symmetry breaking caused by the existence of Dirac spinor fields with the exact Lorentz symmetry group [2, 9]. Describing spinor fields in terms of the composite bundle (5), we get their Lagrangian (26) in the presence of a general linear connection which is invariant under general covariant transformations [1, 9]. Classical gauge fields also are considered in gauge theory on gauge-natural bundles [7] and in Stelle – West gravitation theory [4].

Let us note however that the symmetry breaking mechanism of Standard Model differs from that we consider here. Matter fields in Standard Model admit a total group of symmetries which are broken because of the existence of a background Higgs vacuum field [6].

References

[1] Giachetta, G., Mangiarotti, L. and Sardanashvily, G. (2009). *Advanced Classical Field Theory* (World Scientific, Singapore).

[2] Ivanenko, D. and Sardanashvily, G. (1983). The gauge treatment of gravity, *Phys. Rep.* **94**, 1.

[3] Kobayashi, S. and Nomizu, K. (1963). *Foundations of Differential Geometry, Vol. 1* (John Wiley, New York).

[4] Leclerc, M. (2006). The Higgs sector of gravitational gauge theories, *Annals Phys.* **321**, 708.

[5] Mackey, G. (1968). *Induced Representations of Groups and Quantum Mechanics* (W.A. Benjamin, New York).

[6] Novaes, S. (2000). Standard model: An introduction, *arXiv*: hep-ph/0001283.

[7] Palese, M. and Winterroth, E. (2013). Higgs fields on gauge-natural bundles, *J. Phys.: Conf. Ser.* **411**, 012025.

[8] Sardanashvily, G. (2006). Geometry of classical Higgs fields, *Int. J. Geom. Methods Mod. Phys.* **3**, 139; *arXiv*: hep-th/0510168.

[9] Sardanashvily, G. (2011). Classical gauge gravitation theory, *Int. J. Geom. Methods Mod. Phys.*, **8**, 1869; *arXiv*: 1110.1176.

[10] Sardanashvily, G., Kurov A. (2013). Theory of classical Higgs fields. I. Matter fields, *arXiv*: 1312.3696.

[11] Sardanashvily, G. (2014). Classical Higgs fields, *Theor. Math. Phys.* **181**, 1598.

[12] Steenrod, N. (1972). *The Topology of Fibre Bundles* (Princeton Univ. Press, Princeton).